

Empower Patient Services start form

Empower Patient Services is administered by Optime Care Specialty Pharmacy, the exclusive pharmacy provider for ORLADEYO™ (berotralstat). That means a single, dedicated point of contact for you and your patient throughout your patient's treatment with ORLADEYO. From initial benefits investigation to delivery and ongoing support, a dedicated care coordinator will help support your patients and work with you through the process.

Complete enrollment form

Return pages 2 to 5

- Prescribing healthcare professional must sign page 3

Provide copies of these documents (if available)

Both sides of insurance card

Both sides of prescription benefit card

Lab results supporting diagnosis of hereditary angioedema (HAE)

- C1-inhibitor quantitative (antigenic)
- C1-inhibitor functional
- C4 quantitative (antigenic)

-
- Other supporting documentation to have available if requested:

- Attack history
- Family history
- Androgen failure or contraindication
- Current HAE medications
- Previous HAE medications
- Letter of medical necessity
- Failure of antihistamines and/or epinephrine
- Notes from most recent office visit

Fax or email completed enrollment form and documents

Empower Patient Services

Fax: 1-844-336-7693

Email: info@EmpowerPS.com

Prescriber information

First name

Last name

Specialty and/or designation

Phone

Email

Site/office name

Street address

City

State

ZIP

NPI

State license No.

Preferred method of contact (select all that apply)

Phone Email Fax

Office phone

Fax

Email

Office information

Preferred contact first name

Preferred contact last name

Insurance information (if available)

Attach both sides of the patient's insurance card and prescription benefit card (if applicable) or fill out the information below.

Primary insurance

Subscriber first name

Subscriber last name

Policy ID

Policy group ID

Rx BIN

Rx PCN

Pharmacist help desk phone

Relationship of subscriber to patient

Other insurance (if applicable)

Subscriber first name

Subscriber last name

Policy ID

Policy group ID

Rx BIN

Rx PCN

Pharmacist help desk phone

Relationship of subscriber to patient

Caregiver (optional)

First name

Last name

Phone

Relationship to patient

Email (optional)

Patient information

First name	Middle initial	Street address		
<input type="text"/>	<input type="text"/>	<input type="text"/>		
Last name	Street address 2 (optional)			
<input type="text"/>	<input type="text"/>			
Date of birth	Gender	City	State	ZIP
<input type="text"/>	M <input type="checkbox"/> F <input type="checkbox"/>	<input type="text"/>	<input type="text"/>	<input type="text"/>
MM	DD	YYYY		
Phone	Alternate phone (optional)	Email		
<input type="text"/>	<input type="text"/>	<input type="text"/>		

ORLADEYO™ (berotralstat) prescription information

Diagnosis:

ICD-10 D84.1

Other (please specify)

Special precautions (eg, allergies):

Dose (select one):

1 (one) capsule (**150 mg**) orally, once daily with food; dispense quantity of 28 capsules, 4-week supply

1 (one) capsule (**110 mg**) orally, once daily with food; dispense quantity of 28 capsules, 4-week supply

Refills:

12

Other

12-week supply (if available):

Yes

Customized dosing directions:

If eligible, I request for my patient to participate in the Quick Start program that will provide free drug during the insurance approval process. The Quick Start program is available to all insured patients ≥12 years of age who are US residents with a confirmed diagnosis of HAE. Eligibility is subject to the terms and conditions of the program. BioCryst reserves the right to rescind, revoke, or amend the program at any time without notice. Contact Empower Patient Services for details.

ORLADEYO prescriber signature

By signing below, I certify that (a) the above therapy is medically necessary and that I will supervise the patient's treatment accordingly; (b) I have received the necessary authorizations, including those required by state law and the Health Insurance Portability and Accountability Act of 1996 (HIPAA), to release the above information and other health and medical information of the patient to the dispensing pharmacy.

Sign only once	}	Dispense as written	_____	_____
			<i>Signature</i>	<i>Date</i>
		OR	<i>Prescriber's full signature is required. No stamps.</i>	
		Substitutions allowed	_____	_____
			<i>Signature</i>	<i>Date</i>
			<i>Prescriber's full signature is required. No stamps.</i>	

New York prescribers—please submit prescription on an original NY state prescription blank.

Please see Important Safety Information on page 6 and accompanying full Prescribing Information. 3

AUTHORIZATION FOR USE AND DISCLOSURE OF PERSONAL HEALTH INFORMATION

If the patient is not present at the time of prescription, the start form can be submitted without patient signature. Empower Patient Services can reach out to the patient once the start form has been submitted.

I authorize Optime Care, Inc. ("Optime") to share my, or my legal dependent's, as applicable, personal health information ("PHI"), including, but not limited to, my medical diagnosis, condition, treatment (including prescription information), health insurance information, financial information, demographic information, and contact information, whether provided to Optime previously or in the future, with BioCryst Pharmaceuticals, Inc. (including its representatives and service providers) ("BioCryst").

I authorize such disclosures so that BioCryst and Optime may use my PHI for the following purposes:

- to provide product support services for ORLADEYO, including, but not limited to, copay assistance, reimbursement support, and other forms of patient assistance
- communicating with me by mail, email, text message, telephone, or other means about my medical condition, treatment, care management, and health insurance
- internal use by BioCryst, including data analysis, to evaluate services and to improve future products and services related to HAE treatment
- to contact me about my interest in participating in market research
- to contact me about participation in a mentor program

I authorize BioCryst and Optime to use my PHI for these purposes and to share my PHI in connection with these purposes, including with my healthcare providers, insurance providers, and pharmacy, and their representatives, in order for them to coordinate my benefits, provide, when applicable, reimbursement support, investigate my insurance coverage, and help with financial assistance for BioCryst products.

I also authorize Optime to share my PHI related to my HAE condition and treatment with the patient support organizations related to HAE ("Support Organizations"), including their representatives and service providers.

I authorize such disclosures so that Support Organizations may use my PHI for the following purposes:

- to provide support related to my medical condition(s), including coordination of benefits and therapy;
- reimbursement support;
- investigating insurance coverage;
- communicating with me by mail, email, text message, telephone, or other means about my medical condition, treatment, care management, and health insurance.

I understand that once my PHI is shared, the information may no longer be protected by federal privacy regulations and could be re-disclosed, but that the intent is to use my PHI only for the purposes listed above. I understand that I do not have to sign this Authorization in order to receive healthcare, payment for healthcare, or to be eligible for healthcare benefits.

This Authorization expires 20 years from the date of my signature below, unless otherwise required by law.

I agree that if I reside in the state of Maryland, this form will be valid for no longer than 1 year from the date signed.

I have the right to cancel this authorization. If I cancel, this means that BioCryst and/or Support Organizations will no longer use or share my PHI, but this will not apply to PHI already used or shared or when it is required by law.

If I reside in California, I also have the right to request that BioCryst and/or Support Organizations delete my PHI, although deletion is not required under certain circumstances. To cancel or request deletion, I must send a written notice to BioCryst. It can be sent by fax or by mail to the address below. If I cancel and request deletion, I know that BioCryst and Support Organizations will no longer be able to assist me with access to ORLADEYO.

I understand that I may revoke this Authorization by sending a written notice of revocation to 4060 Wedgeway Court, Earth City, MO 63045. I understand that if I do revoke the Authorization, that will not invalidate any uses or disclosures of my PHI made in reliance on the Authorization prior to the receipt by Optime, BioCryst, and Support Organizations of my notice of revocation.

I understand that I am entitled to receive a copy of this Authorization over the time it is valid.

I certify that I am at least eighteen (18) years of age.

By signing below, I certify that I have read and agree to the above

Patient signature

Patient full signature is required. No stamps.

Date

Patient printed name

If being signed by authorized representative, please sign below and describe your relationship to patient

Signature

Patient's legally authorized representative signature (if applicable)

Date

First name

Last name

Relationship to the patient

INDICATION

ORLADEYO™ (berotralstat) is a plasma kallikrein inhibitor indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adults and pediatric patients 12 years and older.

Limitations of use

The safety and effectiveness of ORLADEYO for the treatment of acute HAE attacks have not been established. ORLADEYO should not be used for the treatment of acute HAE attacks. Additional doses or dosages of ORLADEYO higher than 150 mg once daily are not recommended due to the potential for QT prolongation.

IMPORTANT SAFETY INFORMATION

An increase in QT prolongation was observed at dosages higher than the recommended 150 mg once-daily dosage and was concentration dependent.

The most common adverse reactions ($\geq 10\%$ and higher than placebo) in patients receiving ORLADEYO were abdominal pain, vomiting, diarrhea, back pain, and gastroesophageal reflux disease.

A reduced dosage of 110 mg taken orally once daily with food is recommended in patients with moderate or severe hepatic impairment (Child-Pugh B or C) and in patients taking chronically administered P-glycoprotein (P-gp) or breast cancer resistance protein (BCRP) inhibitors (eg, cyclosporine).

Berotralstat is a substrate of P-gp and BCRP. P-gp inducers (eg, rifampin, St. John's wort) may decrease berotralstat plasma concentration, leading to reduced efficacy of ORLADEYO. The use of P-gp inducers is not recommended with ORLADEYO.

ORLADEYO at a dose of 150 mg is a moderate inhibitor of CYP2D6 and CYP3A4. For concomitant medications with a narrow therapeutic index that are predominantly metabolized by CYP2D6 or CYP3A4, appropriate monitoring and dose titration is recommended. ORLADEYO at a dose of 300 mg is a P-gp inhibitor. Appropriate monitoring and dose titration is recommended for P-gp substrates (eg, digoxin) when coadministering with ORLADEYO.

The safety and effectiveness of ORLADEYO in pediatric patients <12 years of age have not been established.

There are insufficient data available to inform drug-related risks with ORLADEYO use in pregnancy. There are no data on the presence of berotralstat in human milk, its effects on the breastfed infant, or its effects on milk production.

To report SUSPECTED ADVERSE REACTIONS, contact BioCryst Pharmaceuticals, Inc. at 1-833-633-2279 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see accompanying full Prescribing Information.

Next steps

1. Submit the start form and supporting documentation via fax or email.
2. Empower Patient Services will confirm receipt of form via email or fax, depending on office preference.
3. Your dedicated care coordinator that is assigned to your office will get in touch with your patient and verify benefits and the health plan process for the patient. He or she will also connect with your office regarding any missing information.
4. If applicable, your care coordinator will assess your patient's eligibility for financial assistance and support programs.
5. Your dedicated care coordinator will work alongside you, your office, and your patient throughout the entire approval process as well as provide ongoing support after initiation of treatment with ORLADEYO.

HIGHLIGHTS OF PRESCRIBING INFORMATION

These highlights do not include all the information needed to use ORLADEYO™ safely and effectively. See full prescribing information for ORLADEYO™.

ORLADEYO™ (berotralstat) capsules, for oral use
Initial U.S. Approval: 2020

INDICATIONS AND USAGE

ORLADEYO is a plasma kallikrein inhibitor indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adults and pediatric patients 12 years and older. (1)

Limitations of Use:

ORLADEYO should not be used for treatment of acute HAE attacks. (1)

DOSAGE AND ADMINISTRATION

• Recommended Dosage: One capsule (150 mg) taken orally once daily with food. (2.1)

See Full Prescribing Information for:

- Dosage adjustment in patients with moderate or severe hepatic impairment. (2.2)
- Dosage adjustment in patients with chronic administration of P-gp or BCRP inhibitors. (2.3)
- Dosage adjustment in patients with persistent gastrointestinal reactions. (2.4)

DOSAGE FORMS AND STRENGTH

Capsules: 150 mg, 110 mg (3)

CONTRAINDICATIONS

None (4)

WARNINGS AND PRECAUTIONS

An increase in QT prolongation can occur at dosages higher than the recommended 150 mg once daily dosage. Additional doses or doses of ORLADEYO higher than 150 mg once daily are not recommended. (5.1)

ADVERSE REACTIONS

Most common adverse reactions (≥10%) are abdominal pain, vomiting, diarrhea, back pain, and gastroesophageal reflux disease. (6.1)

To report SUSPECTED ADVERSE REACTIONS, contact BioCryst Pharmaceuticals, Inc. at 1-833-633-2279 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DRUG INTERACTIONS

P-gp or BCRP inhibitors – Reduce ORLADEYO dosage when co-administered. (7.1, 12.3)

P-gp inducers – Avoid use with ORLADEYO. (7.1)

CYP2D6, CYP3A4 or P-gp Substrates: Appropriately monitor or dose titrate narrow therapeutic index drugs that are predominantly metabolized by CYP2D6, CYP3A4 or are P-gp substrates when co-administered with ORLADEYO. (7.2, 12.3)

See 17 for PATIENT COUNSELING INFORMATION and FDA-approved patient labeling.

Revised: 12/2020

FULL PRESCRIBING INFORMATION: CONTENTS*

1 INDICATIONS AND USAGE

2 DOSAGE AND ADMINISTRATION

- 2.1 Recommended Dosage
- 2.2 Recommended Dosage in Patients with Hepatic Impairment
- 2.3 Recommended Dosage for Concomitant Use with P-gp or BCRP Inhibitors
- 2.4 Dosage Adjustment in Patients with Persistent GI Reactions

3 DOSAGE FORMS AND STRENGTHS

4 CONTRAINDICATIONS

5 WARNINGS AND PRECAUTIONS

- 5.1 Risk of QT Prolongation with Higher-Than-Recommended Dosages

6 ADVERSE REACTIONS

- 6.1 Clinical Trials Experience

7 DRUG INTERACTIONS

- 7.1 Potential for Other Drugs to Affect ORLADEYO
- 7.2 Potential for ORLADEYO to Affect Other Drugs

8 USE IN SPECIFIC POPULATIONS

- 8.1 Pregnancy
- 8.2 Lactation
- 8.4 Pediatric Use
- 8.5 Geriatric Use
- 8.6 Renal Impairment
- 8.7 Hepatic Impairment

11 DESCRIPTION

12 CLINICAL PHARMACOLOGY

- 12.1 Mechanism of Action
- 12.2 Pharmacodynamics
- 12.3 Pharmacokinetics

13 NONCLINICAL TOXICOLOGY

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

14 CLINICAL STUDIES

16 HOW SUPPLIED/STORAGE AND HANDLING

17 PATIENT COUNSELING INFORMATION

* Sections or subsections omitted from the full prescribing information are not listed

FULL PRESCRIBING INFORMATION

1 INDICATIONS AND USAGE

ORLADEYO™ is indicated for prophylaxis to prevent attacks of hereditary angioedema (HAE) in adults and pediatric patients 12 years of age and older.

Limitations of Use:

The safety and effectiveness of ORLADEYO for the treatment of acute HAE attacks have not been established. ORLADEYO should not be used for treatment of acute HAE attacks. Additional doses or doses of ORLADEYO higher than 150 mg once daily are not recommended due to the potential for QT prolongation [see *Warnings and Precautions (5.1)*].

2 DOSAGE AND ADMINISTRATION

2.1 Recommended Dosage

The recommended dosage of ORLADEYO is one 150 mg capsule taken orally once daily with food.

2.2 Recommended Dosage in Patients with Hepatic Impairment

No dosage adjustment of ORLADEYO is recommended for patients with mild hepatic impairment (Child-Pugh Class A) [see *Use in Specific Populations (8.7)* and *Clinical Pharmacology (12.3)*].

In patients with moderate or severe hepatic impairment (Child-Pugh B or C), the recommended dosage of ORLADEYO is one 110 mg capsule taken orally once daily with food [see *Use in Specific Populations (8.7)* and *Clinical Pharmacology (12.3)*].

2.3 Recommended Dosage for Concomitant Use with P-gp or BCRP Inhibitors

In patients with chronic administration of P-gp or BCRP inhibitors (e.g., cyclosporine), the recommended dosage of ORLADEYO is one 110 mg capsule taken orally once daily with food [see *Drug Interactions (7.1)* and *Clinical Pharmacology (12.3)*].

2.4 Dosage Adjustment in Patients with Persistent GI Reactions

Gastrointestinal (GI) reactions may occur in patients receiving ORLADEYO [see *Adverse Reactions (6.1)*]. If GI events persist, a reduced dose of 110 mg once daily with food may be considered.

3 DOSAGE FORMS AND STRENGTHS

Capsules:

- 150 mg: a white opaque body with a black imprint “150” and a light blue opaque cap with a black imprint “BCX”.
- 110 mg: light blue opaque capsules with a white imprint “110” on body and a white imprint “BCX” on cap.

4 CONTRAINDICATIONS

None

5 WARNINGS AND PRECAUTIONS

5.1 Risk of QT Prolongation with Higher-Than-Recommended Dosages

ORLADEYO should not be used for treatment of acute attacks of HAE. Additional doses or doses of ORLADEYO higher than 150 mg once daily are not recommended. An increase in QT was observed at dosages higher than the recommended 150 mg once daily dosage and was concentration dependent [see *Clinical Pharmacology (12.2)*].

6 ADVERSE REACTIONS

The following clinically significant adverse reaction is described elsewhere in the labeling:

- QT Prolongation [see *Warnings and Precautions (5.1)*]

6.1 Clinical Trials Experience

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

The safety of ORLADEYO is primarily based on 24-week (Part 1) data from a 3-part, double-blind, parallel-group, and placebo-controlled study (Trial 1) in 120 patients with Type I or II HAE randomized and dosed with either ORLADEYO 110 mg, 150 mg or placebo, once daily with food. After Week 24, patients who continued in the study received active treatment through 48 weeks.

In Trial 1, a total of 81 patients aged 12 years and older with HAE received at least one dose of ORLADEYO in Part 1. Overall, 66% of patients were female and 93% of patients were Caucasian with a mean age of 41.6 years. The proportion of patients who discontinued study drug prematurely due to adverse reactions was 7% and 3% for patients treated with 110 mg and 150 mg ORLADEYO, respectively, and 3% for placebo-treated patients. No deaths occurred in the trial.

The safety profile of ORLADEYO was generally similar across all subgroups of patients, including analysis by age, sex, and geographic region.

Table 1 shows adverse reactions occurring in $\geq 10\%$ of patients in any ORLADEYO treatment group that also occurred at a higher rate than in the placebo treatment group in Trial 1.

Table 1: Adverse Reactions Observed in $\geq 10\%$ of Patients in any ORLADEYO Treatment Group (Trial 1)

Adverse Reaction	Placebo (N=39)	ORLADEYO		
		110 mg (N=41)	150 mg (N=40)	Total (N=81)
	n (%)	n (%)	n (%)	n (%)
Abdominal Pain*	4 (10)	4 (10)	9 (23)	13 (16)
Vomiting	1 (3)	4 (10)	6 (15)	10 (12)
Diarrhea†	0	4 (10)	6 (15)	10 (12)
Back Pain	1 (3)	1 (2)	4 (10)	5 (6)
Gastroesophageal Reflux Disease	0	4 (10)	2 (5)	6 (7)

* includes Abdominal pain, Abdominal discomfort, Abdominal pain upper, and Abdominal tenderness

† includes Diarrhea and Frequent bowel movements

Gastrointestinal reactions, including abdominal pain, vomiting, and diarrhea occurred more frequently in patients receiving ORLADEYO 150 mg versus ORLADEYO 110 mg or placebo. These reactions generally occurred early after initiation of treatment with ORLADEYO, became less frequent with time, and typically self-resolved. No patients in the ORLADEYO 150 mg dose group and 1 patient in the ORLADEYO 110 mg dose group discontinued treatment due to a gastrointestinal adverse reaction.

Less Common Adverse Reactions

Other adverse reactions that occurred in Part 1 of Trial 1 with an incidence between 5% and <10% at a higher incidence in ORLADEYO-treated patients compared to placebo included headache (9% versus 5%), fatigue (6% versus 3%), and flatulence (6% versus 3%).

A maculopapular drug rash was reported in less than 1% of patients treated with ORLADEYO. The rash resolved, including in subjects who continued dosing.

Safety data are also available from 227 patients enrolled in an ongoing, open-label, long-term safety study (Trial 2) who received ORLADEYO 110 mg (N=100) or 150 mg (N=127) once daily with food and are consistent with the 24-week controlled safety data from Trial 1 (Part 1).

Laboratory Abnormalities

Transaminase elevations

In Part 1 of Trial 1, a single 150 mg ORLADEYO-treated patient discontinued treatment due to asymptomatic elevated transaminases (ALT >8x the upper limit of normal [ULN] and AST >3x ULN). Total bilirubin was normal. No subject receiving 110 mg or placebo developed transaminase levels >3x ULN. In addition to this patient, 2 ORLADEYO-treated patients developed laboratory-related hepatic adverse events compared to 1 placebo-treated patient. No patient reported serious adverse reactions of elevated transaminases.

7 DRUG INTERACTIONS

This section describes clinically relevant drug interactions with ORLADEYO. Drug interaction studies are described elsewhere in the labeling [*see Clinical Pharmacology (12.3)*].

7.1 Potential for Other Drugs to Affect ORLADEYO

P-gp or BCRP inhibitors

ORLADEYO is a P-gp and BCRP substrate. A dose of 110 mg ORLADEYO is recommended for patients with chronic administration of P-gp or BCRP inhibitors (e.g., cyclosporine) [*see Clinical Pharmacology (12.3)*].

P-gp Inducers

Berotrastat is a substrate of P-gp and BCRP. P-gp inducers (e.g., rifampin, St. John's wort) may decrease berotrastat plasma concentration, leading to reduced efficacy of ORLADEYO. The use of P-gp inducers is not recommended with ORLADEYO.

7.2 Potential for ORLADEYO to Affect Other Drugs

CYP2D6 and CYP3A4 Substrates

ORLADEYO at a dose of 150 mg is a moderate inhibitor of CYP2D6 and CYP3A4. For concomitant medications with a narrow therapeutic index that are predominantly metabolized by CYP2D6 (e.g., thioridazine, pimozide) or CYP3A4 (e.g., cyclosporine, fentanyl), appropriate monitoring and dose titration is recommended [*see Clinical Pharmacology (12.3)*].

P-gp Substrates

ORLADEYO at a dose of 300 mg is a P-gp inhibitor. Appropriate monitoring and dose titration is recommended for P-gp substrates (e.g. digoxin) when co-administering with ORLADEYO [*see Clinical Pharmacology (12.3)*].

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Risk Summary

There are insufficient data in pregnant women available to inform drug-related risks with ORLADEYO use in pregnancy. Based on animal reproduction studies, no evidence of structural alterations was observed when berotralstat was administered orally to pregnant rats and rabbits during organogenesis at doses up to approximately 10 and 2 times, respectively, the maximum recommended human daily dose (MRHDD) in adults on an AUC basis (*see Data*).

The background risk of major birth defects and miscarriage for the indicated population is unknown. In the U.S. general population, the estimated background risk of major birth defects and miscarriage in clinically recognized pregnancies is 2-4% and 15-20%, respectively.

Data

Animal Data

In animal reproduction studies, oral administration of berotralstat to pregnant rats and rabbits during the period of organogenesis did not cause fetal structural alterations. The berotralstat dose in rats and rabbits was up to approximately 10 and 2 times, respectively, the MRHDD in adults (on an AUC basis at maternal doses of 75 and 100 mg/kg/day, respectively). In a pre- and postnatal development study in rats, oral administration of berotralstat to pregnant rats during the period of organogenesis and until delivery at doses up to 45 mg/kg/day (approximately 2 times of the MRHDD on a mg/m² basis) did not cause fetal structural alterations either. Berotralstat concentrations in the fetal blood were approximately 5-11% of the maternal blood.

8.2 Lactation

Risk Summary

There are no data on the presence of berotralstat in human milk, its effects on the breastfed infant, or its effects on milk production. However, when a drug is present in animal milk, it is likely that the drug will be present in human milk. Low levels of berotralstat were detected in the plasma of rat pups when dams were dosed with the drug orally during the lactation period. The berotralstat concentration in the pup plasma was approximately 2% of the maternal plasma (*see Data*).

The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for ORLADEYO and any potential adverse effects on the breastfed infant from ORLADEYO or from the underlying maternal condition.

Data

Animal Data

In the pre- and post-natal development study in rats, berotralstat was administered to dams during the pregnancy and lactation periods at doses up to 45 mg/kg/day (approximately 2 times of the MRHDD on a mg/m² basis). Berotralstat was detected in the plasma of pups during the lactation period. The berotralstat concentration in the pup plasma was approximately 2% of the maternal plasma. Both dams and pups at 45 mg/kg/day showed statistically significant decreases in body weight gain ($p < 0.05$). No treatment-related effects were observed at 25 mg/kg/day (approximately equal to the MRHDD on a mg/m² basis).

8.4 Pediatric Use

The safety and effectiveness of ORLADEYO for prophylaxis to prevent attacks of hereditary angioedema have been established in pediatric patients aged 12 and older. Use of ORLADEYO in this population is supported by evidence from an adequate and well-controlled study (Trial 1) that included adults and a total of 6 adolescent patients aged 12 to <18 years of age. The safety profile

and attack rate on study were similar to those observed in adults [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)* and *Clinical Studies (14)*]. An additional 10 adolescent patients aged 12 to <18 years were enrolled in the open-label study (Trial 2).

The safety and effectiveness of ORLADEYO in pediatric patients < 12 years of age have not been established.

8.5 Geriatric Use

The safety and effectiveness of ORLADEYO were evaluated in a subgroup of patients (N=9) aged \geq 65 years in Trial 1. Results of the subgroup analysis by age were consistent with overall study results. The safety profile from an additional 5 elderly patients aged \geq 65 years enrolled in the open-label, long-term safety study (Trial 2) was consistent with data from Trial 1 [see *Adverse Reactions (6.1)*, *Clinical Pharmacology (12.3)* and *Clinical Studies (14)*].

8.6 Renal Impairment

No dosage adjustment of ORLADEYO is recommended for patients with mild, moderate or severe renal impairment [see *Clinical Pharmacology (12.3)*].

ORLADEYO has not been studied in patients with End-Stage Renal Disease ($CL_{CR} < 15$ mL/min or $eGFR < 15$ mL/min/ 1.73 m² or patients requiring hemodialysis), and, therefore is not recommended for use in these patient populations [see *Clinical Pharmacology (12.3)*].


8.7 Hepatic Impairment

No dosage adjustment of ORLADEYO is recommended for patients with mild hepatic impairment (Child-Pugh Class A) [see *Clinical Pharmacology (12.3)*].

In patients with moderate or severe hepatic impairment (Child-Pugh B or C), the recommended dose of ORLADEYO is 110 mg once daily with food [see *Dosage and Administration (2.2)* and *Clinical Pharmacology (12.3)*].

11 DESCRIPTION

ORLADEYO (berotralstat) capsules is a plasma kallikrein inhibitor. Berotralstat is presented as the dihydrochloride salt with the chemical name 1-[3-(aminomethyl)phenyl]-N-(5-((R)-3-cyanophenyl)[(cyclopropylmethyl)amino]methyl)-2-fluorophenyl)-3-(trifluoromethyl)-1H-pyrazole-5-carboxamide dihydrochloride. The chemical structure is:



Berotralstat dihydrochloride is a white to off-white powder that is soluble in water at $pH \leq 4$. The molecular formula is $C_{30}H_{26}F_4N_6O \cdot 2HCl$ and the molecular weight is 635.49 (dihydrochloride).

ORLADEYO is supplied as 150 mg (equivalent to 169.4 mg berotralstat dihydrochloride) and 110 mg (equivalent to 124.2 mg berotralstat dihydrochloride) hard gelatin capsules for oral administration. Each capsule contains the active ingredient berotralstat dihydrochloride and the inactive ingredients colloidal silicon dioxide, crospovidone, magnesium stearate, and pregelatinized starch.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Berotrastat is a plasma kallikrein inhibitor that binds to plasma kallikrein and inhibits its proteolytic activity. Plasma kallikrein is a protease that cleaves high-molecular-weight-kininogen (HMWK) to generate cleaved HMWK (cHMWK) and bradykinin, a potent vasodilator that increases vascular permeability resulting in swelling and pain associated with HAE. In patients with HAE due to C1-inhibitor (C1-INH) deficiency or dysfunction, normal regulation of plasma kallikrein activity is not present, which leads to uncontrolled increases in plasma kallikrein activity and results in angioedema attacks. Berotrastat decreases plasma kallikrein activity to control excess bradykinin generation in patients with HAE.

12.2 Pharmacodynamics

Concentration-dependent inhibition of plasma kallikrein, measured as a reduction from baseline of specific enzyme activity, was demonstrated after oral administration of ORLADEYO once daily in patients with HAE.

Cardiac Electrophysiology

At the recommended dose of 150 mg once daily, ORLADEYO does not prolong the QT interval to any clinically relevant extent. At 3-times the recommended dose, the mean (upper 90% confidence interval) increase in QTcF was 15.9 msec (23.5 msec). The observed increase in QTcF was concentration-dependent.

12.3 Pharmacokinetics

Following oral administration of berotrastat 150 mg once daily, the steady state C_{max} and area under the curve over the dosing interval (AUC_{tau}) are 158 ng/mL (range: 110 to 234 ng/mL) and 2770 ng*hr/mL (range: 1880 to 3790 ng*hr/mL), respectively. Following oral administration of berotrastat 110 mg once daily, the steady-state C_{max} and AUC_{tau} are 97.8 ng/mL (range: 63 to 235 ng/mL) and 1600 ng*hr/mL (range: 950 to 4170 ng*hr/mL), respectively.

Berotrastat exposure (C_{max} and AUC) increases greater than proportionally with dose and steady state is reached by days 6 to 12. After once-daily administration, exposure of berotrastat at steady state is approximately 5 times that after a single dose.

The pharmacokinetics of berotrastat are similar between healthy adult subjects and in patients with HAE.

Absorption

The median time to maximum plasma concentration (T_{max}) of berotrastat when administered with food is 5 hours (range: 1 to 8 hours).

Effect of Food

No differences in the C_{max} and AUC of berotrastat were observed following administration with a high-fat meal, however the median T_{max} was delayed by 3 hours, from 2 hours (fasted) to 5 hours (fed).

Distribution

Plasma protein binding is approximately 99%. After a single dose of radiolabeled berotrastat 300 mg, the blood to plasma ratio was approximately 0.92.

Elimination

The median elimination half-life of berotrastat was approximately 93 hours (range: 39 to 152 hours).

Metabolism

Berotrastat is metabolized by CYP2D6 and by CYP3A4 with low turnover in vitro. After a single oral radiolabeled berotrastat 300 mg dose, berotrastat represented 34% of the total plasma radioactivity, with 8 metabolites, each accounting for between 1.8 and 7.8% of the total radioactivity.

Excretion

After a single oral radiolabeled berotrastat 300 mg dose, approximately 9% was excreted in urine (3.4% unchanged; range 1.8 to 4.7%) and 79% was excreted in feces.

Specific Populations

Body weight, age, gender, and race did not have a clinically meaningful influence on the systemic exposure of berotrastat.

Geriatric Patients

Based on the population pharmacokinetic analyses that included elderly patients (≥ 65 to 74 years, N=25), age does not have a clinically meaningful impact on the systemic exposure of berotrastat [see *Use in Specific Populations (8.5)*].

Pediatric Patients

Based on population pharmacokinetic analyses that included pediatric patients 12 to <18 years of age, exposure at steady state following oral administration of berotrastat 150 mg once daily was approximately 20% higher compared to adults. The higher exposure in adolescents is not considered to be clinically meaningful.

Patients with Renal Impairment

The pharmacokinetics of a single 200 mg oral dose of berotrastat were studied in subjects with severe renal impairment (CL_{CR} less than 30 mL/min). When compared to a concurrent cohort with normal renal function (CL_{CR} greater than 90 mL/min), no clinically relevant differences were observed; C_{max} was increased by 47%, while AUC_{0-last} was increased by 14% [see *Use in Specific Populations (8.6)*].

The pharmacokinetics of berotrastat has not been studied in patients with End-Stage Renal Disease (CL_{CR} less than 15 mL/min or eGFR less than 15 mL/min/1.73 m² or patients requiring hemodialysis).

Patients with Hepatic Impairment

The pharmacokinetics of a single 150 mg oral dose of berotrastat were studied in subjects with mild, moderate and severe hepatic function (Child-Pugh Class A, B, and C, respectively). The pharmacokinetics of berotrastat were unchanged in subjects with mild hepatic impairment compared to subjects with normal hepatic function. In subjects with moderate hepatic impairment; C_{max} was increased by 77%, while AUC_{0-inf} was increased by 78%. In subjects with severe hepatic impairment, C_{max} was increased by 27%, while AUC_{0-last} was decreased by 5%. The median half-life of berotrastat was increased by 37% and 22% in patients with moderate and severe hepatic impairment, respectively, in comparison to healthy subjects. The percent of unbound berotrastat increased 2-fold from a mean of 1.2% in healthy subjects to a mean of 2.4% in subjects with severe hepatic impairment [see *Use in Specific Populations (8.7)*].

Drug Interaction Studies

Effect of Other Drugs on the Pharmacokinetics of ORLADEYO

Berotrastat is a P-gp and BCRP substrate. Cyclosporine, a P-gp and BCRP inhibitor, increased berotrastat C_{max} by 25%, AUC_{0-last} by 55%, and AUC_{0-inf} by 69% [see *Drug Interactions (7.1)*].


Effect of ORLADEYO on the Pharmacokinetics of Other Drugs

Bertralstat 150 mg once daily is a moderate inhibitor of CYP2D6 and CYP3A4, and a weak inhibitor of CYP2C9 and CYP2C19.

Bertralstat at 300 mg dose is an inhibitor of P-gp and is not an inhibitor of BCRP (rosuvastatin exposure was decreased by approximately 20%).

The effect of bertralstat on the pharmacokinetics of other drugs are presented in Figure 1 [see Drug Interactions (7.2)].

Figure 1: Effect of ORLADEYO on Concomitant Medications



13 NONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Carcinogenesis

Carcinogenicity of bertralstat was evaluated in a 2-year study in Wistar rats and a 26-week study in Tg.rasH2 transgenic mice. The bertralstat doses (oral gavage) were up to 20 and 50 mg/kg/day in rats and mice (approximately 5 and 10 times the MRHDD on a plasma AUC basis, respectively). No evidence of tumorigenicity was observed in either species.

Mutagenesis

Berotrastat tested negative in the *in vitro* bacterial reverse mutation assay (Ames test), the *in vitro* chromosomal aberration assay in human peripheral blood lymphocytes, and the *in vivo* rat micronucleus assay.

Impairment of Fertility

In a fertility study in rats, berotrastat at oral doses up to 45 mg/kg/day (approximately 2 times the MRHDD on a mg/m² basis) showed no effect on fertility in males or females.

14 CLINICAL STUDIES

Trial 1 (NCT3485911)

The efficacy of ORLADEYO for the prevention of angioedema attacks in patients 12 years of age and older with Type I or II HAE was demonstrated in Part 1 of a multicenter, randomized, double-blind, placebo-controlled, parallel-group study (Trial 1).

The study included 120 adult and adolescent patients who experienced at least two investigator-confirmed attacks within the first 8 weeks of the run-in period and took at least one dose of study treatment. Patients were randomized into 1 of 3 parallel treatment arms, stratified by baseline attack rate, in a 1:1:1 ratio (berotrastat 110 mg, berotrastat 150 mg, or placebo by oral administration once daily, with food) for the 24-week treatment period (Part 1).

Patients discontinued other prophylactic HAE medications prior to entering the study; however, all patients were allowed to use rescue medications for treatment of breakthrough HAE attacks.

A history of laryngeal angioedema attacks was reported in 74% of patients and 75% reported prior use of long-term prophylaxis. The median attack rate during the prospective run-in period (baseline attack rate) was 2.9/month. Seventy percent of patients enrolled had a baseline attack rate of ≥ 2 attacks/month.

ORLADEYO 150 mg and 110 mg produced statistically significant reductions in the rate of HAE attacks compared to placebo for the primary endpoint in the Intent-to-Treat (ITT) population as shown in Table 2. The percent reductions in HAE attack rate were greater with ORLADEYO 150 mg and 110 mg relative to placebo regardless of attack rate during the run-in period.

Table 2. Primary Efficacy Endpoint (Trial 1): Reduction in HAE Attack Rate- ITT Population

Outcome	ORLADEYO		Placebo
	110 mg QD	150 mg QD	
	N = 41	N = 40	N = 40*
HAE Attack rate, rate per 28 days†	1.65	1.31	2.35
% Rate Reduction ‡ (95% CI)	30.0% (4.6, 48.7)	44.2% (23.0, 59.5)	
p-value	0.024	<0.001	


* One patient in the ITT analysis was randomized to placebo but was not treated.

† Statistical analysis based on a negative binomial regression model; number of attacks included as dependent variable, treatment included as fixed effect, baseline attack rate included as covariate, and logarithm of duration on treatment included as offset variable.

‡ Percent reduction relative to placebo.

Reductions in attack rates were observed in the first month of treatment with ORLADEYO 150 mg and 110 mg and were sustained through 24 weeks as shown in Figure 2.

Figure 2. Mean (+/- SEM) HAE Attack Rate/month Through 24 Weeks (Trial 1)- ITT Population



Pre-defined exploratory endpoints included the proportion of responders to study drug, defined as at least a 50% relative reduction in HAE attacks during treatment compared with the baseline attack rate; 58% of patients receiving 150 mg ORLADEYO and 51% of patients receiving 110 mg ORLADEYO had a $\geq 50\%$ reduction in their HAE attack rates compared to baseline versus 25% of placebo patients. In post-hoc analyses, 50% and 23% of patients receiving 150 mg ORLADEYO, and 27% and 10% of patients receiving 110 mg ORLADEYO, had a $\geq 70\%$ or $\geq 90\%$ reduction in their HAE attack rates compared to baseline versus 15% and 8% of placebo patients, respectively. The rate of attacks rated as moderate or severe was reduced by 40% and 10% in patients receiving 150 mg ORLADEYO and 110 mg ORLADEYO, respectively, versus placebo.

16 HOW SUPPLIED/STORAGE AND HANDLING

ORLADEYO (berotralstat) capsules:

- 150 mg: a white opaque body with a black imprint “150” and a light blue opaque cap with a black imprint “BCX”. NDC 72769-101-01.
- 110 mg: light blue opaque capsules with a white imprint “110” on body and a white imprint “BCX” on cap. NDC 72769-102-01.
- A 28-day supply of ORLADEYO is provided in a carton containing four child-resistant shellpaks, each containing a 7-capsule blister card.
- Each carton contains a tamper evident seal.
- Do not use if tamper evident seal is broken or missing.

Store at 20°C to 25°C (68°F to 77°F). Excursions permitted between 15°C and 30°C (59°F to 86°F) [see USP Controlled Room Temperature].

17 PATIENT COUNSELING INFORMATION

Advise the patient to read the FDA-approved patient labeling (Patient Information).

Inform patients of the risks and benefits of ORLADEYO before prescribing or administering to the patient.

Drug Interactions

Advise patients that ORLADEYO may interact with other drugs [see *Drug Interactions (7) and Clinical Pharmacology (12.3)*]. Advise patients to report to their healthcare provider the use of any other prescription or nonprescription medication or herbal products.

Not for Acute Treatment of HAE Attacks

Advise patients to take their usual rescue medication to treat an acute attack of HAE. Inform patients that the safety and effectiveness of ORLADEYO has not been established as an acute treatment for HAE attacks. Advise patients that they should not take daily doses higher than 150 mg once daily or additional doses of ORLADEYO to treat an acute attack of HAE due to risk of QT prolongation [see *Limitations of Use (1) and Warnings and Precautions (5.1)*].

For more information, visit www.ORLADEYO.com

ORLADEYO™ is a trademark of BioCryst Pharmaceuticals, Inc.

Manufactured for:
BioCryst Pharmaceuticals, Inc.
Durham, NC 27703

214094-BC-000

PATIENT INFORMATION
ORLADEYO™ (or-luh-DAY-oh)
(berotralstat)
capsules, for oral use

What is ORLADEYO?

- ORLADEYO is a prescription medicine used to prevent attacks of Hereditary Angioedema (HAE) in adults and children 12 years of age and older.
- ORLADEYO is not used to treat an acute HAE attack.
- Do not take more than one capsule of ORLADEYO a day because extra doses can cause heart rhythm problems.
- It is not known if ORLADEYO is safe and effective to treat an acute HAE attack.
- It is not known if ORLADEYO is safe and effective in children under 12 years of age.

Before you take ORLADEYO, tell your healthcare provider about all of your medical conditions, including if you:

- have liver problems or are on kidney dialysis.
- are pregnant or planning to become pregnant. It is not known if ORLADEYO can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if ORLADEYO passes into your breastmilk. Talk to your healthcare provider about the best way to feed your baby while taking ORLADEYO.

Tell your healthcare provider about all of the medicines you take, including other medicines for HAE, prescription and over-the-counter medicines, vitamins, and herbal supplements.

Taking ORLADEYO with certain other medicines may affect the way other medicines work and other medicines may affect how ORLADEYO works.

Know the medicines you take. Keep a list of them to show your healthcare provider and pharmacist when you get a new medicine.

How should I take ORLADEYO?

- Take ORLADEYO exactly as your healthcare provider tells you to take it.
- Take 1 capsule, by mouth, 1 time every day with food.

What are the possible side effects of ORLADEYO?

Taking more than one capsule of ORLADEYO a day may cause serious side effects, including:

- **heart rhythm problems.** A heart rhythm problem called QT prolongation can happen in people who take more than one capsule of ORLADEYO a day. This condition can cause an abnormal heart beat. Do not take more than one capsule of ORLADEYO a day.

The most common side effects of ORLADEYO include:

- abdominal pain
- vomiting
- diarrhea
- back pain
- heartburn

Less common side effects include increases in liver function tests. Rarely, some patients had a brief, itchy rash.

These are not all of the possible side effects of ORLADEYO. For more information, ask your healthcare provider or pharmacist.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

How should I store ORLADEYO?

- Store ORLADEYO at room temperature between 68°F to 77°F (20°C to 25°C).
- Each carton contains a tamper evident seal. Do not use ORLADEYO if the tamper evident seal is broken or missing.

Keep ORLADEYO and all medicines out of the reach of children.

General information about the safe and effective use of ORLADEYO.

Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet. Do not use ORLADEYO for a condition for which it was not prescribed. Do not give ORLADEYO to other people, even if they have the same symptoms that you have. It may harm them. You can ask your pharmacist or healthcare provider for information about ORLADEYO that is written for health professionals.

What are the ingredients in ORLADEYO?

Active ingredient: berotralstat dihydrochloride

Inactive ingredients: colloidal silicon dioxide, crospovidone, magnesium stearate, and pregelatinized starch

Manufactured for: BioCryst Pharmaceuticals, Inc., Durham, NC 27703

ORLADEYO™ and the BioCryst Logo are trademarks or registered trademarks of BioCryst Pharmaceuticals, Inc.

© 2020 BioCryst. All rights reserved.

For more information, visit www.ORLADEYO.com or call 1-833-633-2279.